



A Review of Phytogenic Feed Additives in Aquaculture: Applications, Benefits, and Challenges

Kazempoor R.¹; Alavinezhad S.S.^{2*}; Pargari M.M.³; Shakeri Y.S.³; Haghighi M.M.³

Received: February 2026

Accepted: May 2026

Abstract

Feed production represents the largest operational cost in aquaculture. Consequently, considerable attention has been directed toward the development of sustainable and cost-effective feed additives derived from natural resources. Among these, phytogenics, including herbs, spices, plant extracts, essential oils, oleoresins, and plant-derived by-products, have emerged as promising functional additives for aquatic animal nutrition. In addition to providing a sustainable use for agricultural and fruit-processing wastes, phytogenic additives offer a wide range of biological activities that may enhance fish and shrimp health and performance.

The beneficial effects of phytogenics are largely attributed to their bioactive secondary metabolites, which can stimulate feed intake, improve digestive enzyme secretion, enhance nutrient utilization, modulate intestinal microbiota, and strengthen immune responses. Numerous studies have reported antioxidant, antimicrobial, antiparasitic, and growth-promoting effects of phytogenic compounds in aquatic animals. These properties have increased interest in their use as natural alternatives to antibiotic growth promoters.

Despite the growing body of evidence supporting the application of phytogenics in aquaculture, their mechanisms of action remain incompletely understood. Further research is required to evaluate their long-term effects on host physiology, gut microbiota, animal welfare, and environmental sustainability. A better understanding of these mechanisms will facilitate the safe and effective incorporation of phytogenic additives into aquafeeds.

This review summarizes current knowledge regarding the use of phytogenic feed additives in aquatic animals, with particular emphasis on their sources, biological functions, benefits, limitations, and future research needs.

Keywords: Phytogenics, phytogenic feed additives, aquaculture, fish, shrimp, immune system, growth performance, gut microbiota

1-Department of Biology, Roudehen Branch, Islamic Azad University, Roudehen, Iran.

2-Department of Aquatic Health and Diseases, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran.

3-Department of Aquatic Health and Diseases, Faculty of Veterinary Medicine, Science and Research Branch, Islamic Azad University, Tehran, Iran.

*Corresponding author: shivaalavinejad@ut.ac.ir

Introduction

Phytogenic feed additives or phytobiotics are among the functional feed additives that have been widely studied in recent years. Phytogenic feed additives are non-destructive bioactive compounds of the environment derived from plants, including herbs, spices, essential oils, and extracts. Glycyrrhizin, liquiritin, glabridin, polysaccharides, flavonoids, alkaloids, anthraquinone, saponin, and azadirachtin are among the active ingredients in these phytogenic compounds (Bhujel *et al.*, 2017). Acids, alcohols, aldehydes, acyl esters, sulfur-containing compounds, coumarins, and homologues of polypropanoid are other studied molecules (Yang *et al.*, 2015). A wide range of phytogenics in farmed aquatic animals have been studied mainly due to their properties including improved taste in feed, increased growth, stimulated immune system, improved intestinal microbiota, antimicrobial, antiparasitic and antiviral activities.

Global aquaculture production (including aquatic plants) is 110.2 million tons, with an initial sales value of 243.5 billion dollars. Total production includes 80.0 million tons of edible fish and 30.1 million tons of aquatic plants (Bharathi *et al.*, 2019). According to the FAO, global aquaculture production is estimated to reach 80 million tons by 2050. The aquatic feed contains 50 to 80 percent of rearing costs, and successful aquaculture depends on a balanced diet and low-cost production. As a result, one of the most important challenges for aquaculture

producers is providing adequate and inexpensive nutrients to achieve the ideal growth rate and non-destructive to the environment. The use of low-protein feeds and low amounts of nutrients lead to inefficient use of nutrients. As a result, increased feed intake increases susceptibility to disease and increases production costs. The production of more waste also causes damage to the environment (Goncalves and Santos, 2015).

The presence of endemic pathogens and pandemic diseases constantly endangers aquaculture farmers around the world to achieve a balanced production system. The prevalence of fish diseases is one of the main limitations for the development of aquaculture (FAO, 2020). In the past, antibiotics were used to prevent these problems. But today, producers are looking for alternative ways due to the increasing awareness of consumers about environmental problems and bacterial resistance to the indiscriminate use of antibiotics against zoonotic pathogens. A significant number of studies have focused on the search for antibiotic alternatives with similar antimicrobial and growth-promoting effects without causing residues, bacterial resistance, and potential side effects for animals. Functional feed additives are an alternative to antibiotics and in addition to nutritional value have health benefits for animals (Korzekwa *et al.*, 2016; Waagbø and Remø, 2020). This study aimed to investigate the effects of phytogenic feed additives on aquatic nutrition as an alternative

strategy for antibiotics to achieve sustainable aquaculture.

Classification of phytogenic compounds

Phytogenic feed additives are classified into three main groups, which include sensory additives, technological additives, and zootechnical additives. Some of these additives have more than one positive effect, but they can not be precisely classified in one of the above groups (Karásková *et al.*, 2015).

Sensory additives

This group of additives usually stimulates the appetite through the effect they have on the taste or color of feed (such as plant pigments). The most commonly used sensory additives are carotenoid pigments, which are sourced from carrots, chlorella, marigolds (*Tagetes erecta* L.), or lutein (Karásková *et al.*, 2015). Astaxanthin is one of the known carotenoids in aquatic animals. The use of astaxanthin improves the color of fish fillets (Teimouri *et al.*, 2013). Other studies on carotenoid pigment include increased redness in goldfish and carp (Hancz *et al.*, 2003), production of vivid colors (including blue and green) in marine invertebrates (Davies, 2009), and increased redness in dwarf gourami (Baron *et al.*, 2008). However, the instability and high cost of natural carotenoids have limited the use of these compounds (Englmaierová *et al.* 2014).

Technological additives

This classification refers to a group of additives that affect the technological aspects of the feed. These additives do not directly affect the nutritional value of the feed, but may indirectly affect the nutritional value of the feed by improving maintenance or hygienic properties, such as mycotoxin contaminants in feed (Pandey *et al.*, 2019).

Zootechnical additives

These additives improve the nutrient status and livestock production. These compounds provide specific nutrients and help to use the nutrients in the diet more effectively (Pandey *et al.*, 2019). These additives include immune regulators, digestive stimulants, and growth stimulants of non-microbial origin, and substances that enhance the quality of animal products. These additives are divided into three subgroups: Immunosuppressive compounds (oligosaccharides, especially beta-glucans that stimulate the activity of cells, enzymes, and other factors involved in immunity), compounds that improve production performance or quality of livestock products (cinnamaldehyde and turmeric) They are effective in improving digestion and absorption of nutrients) and compounds that reduce oxidative stress (polyphenols, carotenoids, betalains, and curcumin, which are effective in reducing the level of free radicals (ROS)) (Karásková *et al.*, 2015). The use of these feed additives in aquaculture can be a powerful solution

to limit nitrogen depletion in the environment by improving feed efficiency, saving nutrients, and using proteins, and helping manage fish health (Goncalves and Santos, 2015). According to reports, Feeding with phytogenic feed additives including AROMEX® ME Plus and FRESTA® F Plus, oregano, anise, and citrus peel essential oils reduced excreted ammonia in feed animals (Veit *et al.*, 2011; Hong *et al.*, 2012).

The role of phytoGenics in aquatic growth performance

In recent years, rations containing fish (FM) and fish oil have been used more selectively due to reduced availability and simultaneous price increases, while many efforts are being made to find alternative protein sources for profitable and sustainable aquaculture (FAO, 2018; Slater *et al.*, 2018). Plant products have been proposed as an alternative to FM in aquatic feed due to their low cost, availability, and high protein (Gatlin *et al.*, 2007; Olsen and Hasan, 2012). But the use of plant products is the cause of problems such as high levels of anti-nutritional factors in plant proteins, lack of essential amino acids, lower digestibility, high levels of insoluble carbohydrates, and high fiber, which reduces the palatability of these plant proteins compared to animal proteins (Hardy, 2010), and leads to reduced growth in fish and shrimp that feed on plant protein (De Francesco *et al.*, 2004; Enterria *et al.*, 2011; Montero *et al.*, 2005; Peres and Oliva-Teles, 2006; Sudaryono And Evans, 1999; Zhu *et al.*,

2013a). The use of dietary supplements has been suggested to compensate for the problems caused by replacing FM with a diet containing phytogenic plant proteins. According to studies, phytogenic compounds improve nutrient retention and growth function in aquatic organisms (Karásková *et al.*, 2015; Encarnação, 2016). Induction of secretion of digestive enzymes is one of the mechanisms of phytoGenics in increasing growth performance which stimulates appetite and thus increases feed intake and conversion efficiency (Wenk 2003; Liu *et al.* 2011). For example, spices such as peppers (such as capsaicin and piperine) and other essential oils containing cinnamaldehyde stimulate amylase production (Steiner and Syed 2015). Phytogenic additives (especially essential oils) increase the taste and smell of feed, increase palatability and thus increase feed consumption (Kroismayr *et al.*, 2006). This is probably due to the antioxidant effects of phytogenic compounds, which help maintain the quality of the diet and prevent the spread of unpleasant odors (Franz *et al.*, 2010; Solà-Oriol *et al.*, 2011). However, there is little evidence to increase the palatability of diets containing phytogenic compounds in fish and shrimp (Giannenas *et al.*, 2012; Volpatti *et al.*, 2013).

The role of phytoGenics in the growth of cold water fish

Giannenas *et al.* (2012) reported that using phytogenic additives including carvacrol and thymol showed no

significant difference in feed intake in the diet of Rainbow trout (*Oncorhynchus mykiss*) compared to control diets. But in fish fed with carvacrol and thymol, feed efficiency increased significantly compared to fish fed with the control diet. In the study of Giannenas *et al.* (2012), feeding on diets fortified with carvacrol and thymol improved feed efficiency. But it did not affect fish weight gain compared to the control group. The study by Peterson *et al.* (2014) has not reported the weight gain of trout fed with the commercial phytogenic supplement Digestarom® P.E.P. MGE.

The role of phytogenics in the growth of warm water fish

Zheng *et al.* (2009) evaluated the effect of carvacrol and thymol separately in combination with each other and combination with oregano essential oil in canal catfish. The results of this study showed that fish that consumed oregano essential oil, and fish that used carvacrol alone or a combination of thymol and carvacrol gained more weight than fish in the control group. In a study by Ceulemans *et al.* (2009) on the nutrition of Nile tilapia (*Oreochromis niloticus*) with a diet containing a mixture of plant extracts and natural emulsifying agents, feed conversion and protein consumption efficiency were more favorable than control diets. The use of sweet potato skin in the diet also improved growth performance and nutritional efficiency in Nile tilapia (Omoregie *et al.*, 2009).

The role of phytogenics in shrimp growth

The research by Goncalves and Santos (2015) is one of the studies on the effect of phytosanitary additives on shrimp growth performance. Based on the results of this study, the use of commercial phytogenic feed additive Digestarom® P.E.P. MGE in the diet containing small amounts of fish increased weight, increased feed conversion ratio, and growth rate compared to shrimp that did not use dietary supplements. In another study, papaya leaves increased protein digestion, feed conversion ratio, and growth in *Penaeus monodon* shrimp larvae (Peñaflorida, 1995). In addition, a significant increase in digestive enzymes (amylase, protease, and lipase) and better feed conversion efficiencies were observed in Post-larvae of *P. monodon* fed with enriched Artemia with ginger (*Zingiber officinale*) (Venkatramalingam *et al.*, 2007).

The role of phytogenics in the improvement of immune system function

Numerous studies have reported the use of phytobiotics in aquaculture to stimulate the activity of immune system components (Rao *et al.*, 2006; Sahu *et al.*, 2007; Ardó *et al.*, 2008). Antioxidant effect of phytogenic mechanisms in improving immune function. The antioxidant activity of phytogenic compounds is due to their ability to destroy free radicals, inhibit reactive oxygen species (ROS) and other oxygen radicals produced in cells and tissues.

However, high concentrations of these compounds have been reported to exhibit peroxidation antagonistic effects (Llana-Ruiz-Cabello *et al.*, 2015). Phytochemical compounds are involved in modulating the expression of proinflammatory and anti-inflammatory cytokines by disrupting NF- κ B signaling pathways mediated by TLR and MAPK (Somensi *et al.*, 2019; Zhou *et al.*, 2014; Wu *et al.*, 2017; Liu *et al.*, 2019). In this regard, several studies have shown that phytochemical compounds including curcumin (Shehzad *et al.*, 2011), caffeic acid phenyl ester (CAPE) from bee propolis (Lee *et al.*, 2010), epicatechin (Bahia *et al.*, 2008), grape seed extract and grape brand (GSGME) (Gessner *et al.*, 2013), Cinnamaldehyde (Wondrak *et al.*, 2010) and Purple Potato Anthocyanins (Hwang *et al.*, 2011) increase the expression or transfer of Nrf2 and reduce or inhibit NF- κ B activation. Plant compounds that alter the Nrf2 and NF- κ B pathways can protect the host against oxidative stress, reduce inflammation, and ultimately improve animal health and growth performance. Lectin-mediated pathways are another mechanism by which phytochemical compounds can improve the immune system (LaFrentz *et al.*, 2012) because lectins can activate complement and are involved in improving innate immunity (Ourth *et al.*, 2007).

The role of phytochemicals in improving the immune system of cold water fish

Numerous studies have been performed on the effect of phytochemical additives on improving the immune system function

in cold-water fish (especially rainbow trout). In the study of Giannenas *et al.* (2012) on the effect of feeding rainbow trout with carvacrol and thymol (although the fish were not sick in the challenges) a significant increase in the activity of glutathione-based enzymes (glutathione reductase, glutathione-S-transferase), an increase in lysozyme levels and catalase activity and a significant decrease in serum nitric oxide (NO) were observed. These results indicate the effect of phytochemical additives on intrinsic safety parameters. In the study of Gulec *et al.* (2013), the use of feed containing vegetable oils (*Thymus vulgaris* and *Foeniculum vulgare*) significantly increased the amount of total protein, albumin, cholesterol, triglycerides, bilirubin, K, Na, Ca, Mg, and decreased plasma glucose and chlorine levels in rainbow trout exposed to *Yersinia ruckeri*.

The role of phytochemicals in improving the immune system of warm water fish

According to the results of Yin *et al.* (2006), the extract of Milkvetch root increased phagocytosis of leukocytes and lysozyme activity in the Nile tilapia. In another study, Asimi and Sahu (2013) investigated the effect of feeding on extracts of thyme (*Thymus vulgaris*), rosemary (*Rosmarinus officinalis*), and fenugreek (*Trigonella foenum graecum*) on Mozambique tilapia (*Oreochromis mossambicus*). They observed increased phagocytic activity, increased white and red blood cells, and increased hematocrit levels. In addition, they reported an increase in plasma lysozyme and

myeloperoxidase activity after infection with *Streptococcus iniae*. In *Labeo rohita*, the root extract of *Withania somnifer* increased Nitro blue tetrazolium levels, phagocytic cell activity, lysozyme activity, and total immunoglobulin levels (Sharma *et al.*, 2010). Also, Indian ginseng root extract (Ashwagandha) stimulated the immune system and reduced stress caused by low pH and iron toxicity in water (Laltlanmawia *et al.*, 2019). Injection of Astragalus polysaccharides (APS) increased IL-1 β mRNA levels in a dose-dependent manner in the renal head and regulated TNF- α transcription in the gills and spleen (at high concentrations of APS). TNF- α mRNA was also significantly increased in the renal head (at low concentrations of APS) and regulated Lysozyme-C mRNA levels in gills (at low concentrations of APS) and spleen (at moderate concentrations of APS) in common carp (*Cyprinus carpio*) (Yuan *et al.*, 2008). According to Nahak and Sahu (2014), the use of basil aerial part extract in feed increases RBC, WBC, serum protein, and hematocrit (at concentrations of 2.5 and 5%) and decreases serum glucose (at 5%) in Walking catfish (*Clarias batrachus*) and this compound acts as a strong immune stimulant.

The role of phytogenics in improving the immune system of shrimp

The increase in survival rate is one of the most important effects of increased immunity due to the use of phytogenic compounds in shrimp. For example, the increase in survival rate in the post-

larvae of *Penaeus indicus* shrimp (Immanuel *et al.*, 2004; Citarasu, 2010) and the improvement in the survival rate of *Litopenaeus vannamei* shrimp have been reported as a result of the use of herbal supplements (Yu *et al.*, 2008). It has also been reported that the mixture of Chinese herbal supplements increases nonspecific immunity in shrimp (Luo, 1997; Chansue *et al.*, 2000). He *et al.* (2017) have reported the increased immunity in *Litopenaeus vannamei* shrimp as a result of using stout camphor extract (Yeh *et al.*, 2009) and a combination of organic acids (citric acid and sorbic acid) and essential oils (thymol and vanillin) as dietary supplements. The study by Kesselring *et al.* (2021) showed that feeding a diet low in FM and phytogenic supplementation increased the total number of hemocytes (THCs) and red blood cells compared to a diet high in fish (FM). According to similar studies, the decreased FM levels typically affect shrimp defense mechanisms and reduce hemocytes (Van de Braak *et al.*, 2002). Also, a low-FM diet with phytogenic supplementation reduced ALT and LDH activity levels compared to the low-FM diet group which shows that the use of phytogenic supplements improves the health status of *L. vannamei* by reducing the risk of cell damage and tissue necrosis.

The role of phytogenics in increasing mucosal immunity

Mucosal safety in aquatic animals has been considered by many researchers in recent years due to its diversity and characteristics. Mucosal-associated

lymphoid tissue (MALT) immunity is divided into Gut-associated lymphoid tissue (GALT), Skin-associated lymphoid tissues (SALT), and Gill-associated lymphoid tissue (GIALT) (Lazado and Caipang, 2014). Mode of action of phytochemicals at the level of the

main mucosal lymphoid tissues in fish is depicted in Figure 1. In addition, the effect of phytochemicals against fish pathogenic organisms is suggested in Figure 2.




Figure 1: Suggested mechanisms of cell activation by the transient receptor potential (TRP) cation channels mediated by phytochemicals' bioactive compounds in mucosal-associated lymphoid tissues (MALTs). Bioactive compounds activate TRP channels leading to intracellular Ca^{2+} increase and non-canonical activation of the TAK complex. In parallel, stimulation by pathogen-associated molecular patterns (PAMPs) may facilitate the activation of TLR and TRP signaling pathways. TLR, toll-like receptors; MyD88, myeloid differentiation primary response 88; TAK, transforming growth factor beta (TGF β) activated kinase; TAB, TGF β activated kinase binding protein; ROS, reactive oxygen species; NF- κ B, nuclear factor kappa-B; IKK, inhibitor of NF- κ B kinase; MAPK, mitogen-activated protein kinase; AP1, activator protein 1; CYP, cytochromes P450; P, phosphorylation (Firmino *et al.*, 2021).




Figure 2: Summary of the proposed mode of response induced by phytogenic bioactive compounds in fish mucosal tissues. The effects against representative types of common pathogens on the most studied mucosal-associated lymphoid tissues in fish so far are represented in the figure. (A) Gills. (B) Gut. (C) Skin. Gilthead seabream (*S. aurata*) was used in the figure as a representative aquaculture-relevant fish (Firmino *et al.*, 2021).

Figure 1 Suggested mechanisms of cell activation by the transient receptor potential (TRP) cation channels mediated by phytogenics' bioactive compounds in mucosal-associated lymphoid tissues (MALTs). Bioactive compounds activate TRP channels leading to intracellular Ca^{2+} increase and non-canonical activation of the TAK complex. In parallel, stimulation by pathogen-associated molecular patterns (PAMPs) may facilitate the activation of TLR and TRP signaling pathways. TLR, toll-like receptors; MyD88, myeloid differentiation primary response 88; TAK, transforming growth factor beta ($\text{TGF}\beta$) activated kinase; TAB, $\text{TGF}\beta$

activated kinase binding protein; ROS, reactive oxygen species; NF- κ B, nuclear factor kappa-B; IKK, inhibitor of NF- κ B kinase; MAPK, mitogen-activated protein kinase; AP1, activator protein 1; CYP, cytochromes P450; P, phosphorylation (Firmino *et al.*, 2021).

The role of phytogenics in increasing skin mucosal immunity

The skin plays a key role as the first line of defense against various pathogens in fish (Hoseinifar *et al.*, 2017). Most studies on the effects of phytogenic additives on skin mucosal immunity have shown an increase in skin mucosal immune markers or increased

antibacterial activity of skin mucosa (Heyman *et al.*, 2015; Adel *et al.*, 2015a; Adel *et al.*, 2015b; Gholamhosseini *et al.*, 2020; Heydari *et al.*, 2020; Kurian *et al.*, 2020; Mirghaed *et al.*, 2020; Mohammadi *et al.*, 2020; Paknejad *et al.*, 2020; Rashidian *et al.*, 2021; Yousefi *et al.*, 2021). Lysozyme, alkaline phosphatase, complement and protease activities, total immunoglobulin, and protein content in fish skin mucosa are markers that have been evaluated and have antibacterial potential in vitro against pathogens. Unfortunately, few studies have attempted to identify the underlying mechanisms that regulate the immune system (Firmino *et al.*, 2021).

The role of phytogenics in increasing gill mucosal immunity

Studies on the effects of phytogenic feed additives on gill structure and function are very limited. Some studies have reported the protective effects of these compounds in reducing histopathological lesions of the gills due to exposure to toxic elements or pathogens (Vickers, 2017; Khafaga *et al.*, 2020; El Euony *et al.*, 2020). According to the results of Yuan *et al.* (2008), injection of astragalus polysaccharides (APS) had no effect on IL-1 β mRNA level in the gills but regulated TNF- α and Lysozyme-C mRNA transcription at high and low doses of APS in the gill of common carp (*Cyprinus carpio*).

The role of phytogenics in increasing gastrointestinal mucosal immunity

The gastrointestinal tract (especially the intestine) is mainly composed of physical, chemical, immunological, and microbiological components. Therefore, it is a selective barrier between the host and its surrounding environment (Yegani and Korver, 2008). As a result, any damage or disruption of its normal function suppresses immune responses and reduces animal productivity (McDevitt *et al.*, 2006). In recent years, the study of phytogenic compounds' effect on mucosal immunity has become necessary due to the importance of the mucosal barrier in defending fish against the invasion of pathogens (Yu *et al.*, 2020).

Studies on the effect of phytogenic substances on the intestine include evaluation of changes in morphoanatomical parameters such as improved integrity, increased length, width, and the number of goblet cells (Abdel-Latif *et al.*, 2020; Heluy *et al.*, 2020; Ferreira *et al.*, 2016; Abdel-Tawwab *et al.*, 2018; Vickers, 2017; Araruna *et al.*, 2020), modulation of intestinal lymphocyte count (Hernández *et al.*, 2016; Valladao *et al.*, 2017), improvement in the level of humoral immune markers such as lysozyme (Suttili *et al.*, 2016), improvement in the activity of antioxidant enzymes (Hoseini and Yousefi, 2019; Reyes-Cerpa *et al.*, 2018), reducing the expression of proinflammatory genes such as *tnfa* and *tnfb* (Zhang *et al.*, 2020) and modulation of intestinal microbiota composition (Zhang *et al.*, 2020; Mousavi *et al.*, 2016; Araruna *et al.*, 2020). The positive effects of phytogenics on the

morphology of small intestinal tissues and villi increase nutrient digestibility (Steiner and Syed, 2015). Other effects of phytonics on the gastrointestinal tract include stimulating gastrointestinal secretions and increasing mucus production as a result of increasing the number of goblet cells in the intestine (Goncalves and Santos, 2015). The bacterial status of the gastrointestinal tract changes under the influence of genetic factors, feeding habits, and the environment (Austin, 2006; Wong and Rawls, 2012; Gómez and Balcázar, 2008). However, the intestinal bacterial community and the metabolites produced by them have significant effects on the host health status by modulating the immune system and affecting metabolism (Bento *et al.* 2013; Lazado and Caipang, 2014). Limited studies have examined the effects of phytonics as feed additives on intestinal microbiota in fish. Previous studies have focused on the effect of these plant substances as dietary ingredients instead of additives and their effect on microbial communities of fish intestines (Ringø *et al.*, 2008; Silva *et al.*, 2011; Reveco *et al.* 2014; Zarkasi *et al.*, 2017). Studies on the microbial flora of the gastrointestinal tract are often limited to identifying lactic acid bacteria. Therefore, evaluation of the feed additives' effects on mucosal tissues using the results of these studies is not complete and acceptable and leads to incorrect conclusions. Also, some studies have not reported any effect of phytonics on mucosal parameters (Valladão *et al.*, 2019; Navarrete *et al.*,

2010; Vickers, 2017; Hernández *et al.*, 2015; Vazirzadeh *et al.*, 2019). The use of phytonic compounds as food additives in non-optimal doses and their negligible effect on the bacterial composition of fish intestines is one of the main causes (Suttili *et al.*, 2018).

Studies on the effect of phytonic food additives on the gastrointestinal mucosa are divided into two main groups: the effect on the microbial flora and the intestinal histomorphology. Modification of intestinal microbial flora has been considered by Hongbai *et al.* (2004) using different Chinese plants in carp, Ran *et al.* (2016) using a commercial mixture of thymol and carvacrol in hybrid tilapia (*O. niloticus* x *O. aureus*), and Giannenas *et al.* (2012) using phytonic supplementation containing thymol and carvacrol in rainbow trout. According to a study by Suttili *et al.* (2016) using gradient gel electrophoresis (DGGE) analysis, the use of American basil (*Ocimum americanum*) essential oils with food did not affect the intestinal microbiome in *Sciaenops ocellatus*. In the study by Navarrete *et al.* (2010) using temperature gradient gel electrophoresis (TGGE) analysis, a diet containing thyme-derived essential oil supplement (*Thymus vulgaris*) showed no effect on the intestinal bacterial composition of rainbow trout. In the study of Bello *et al.* (2012), an increase in the height and width of the villi and the depth of crypt was observed in *Clarias gariepinus* fed with onion residues and walnut leaf. Also, according to Goncalves and Santos (2015), the use of commercial

phytogenic additives Digestarom® P.E.P. MGE in a low-fish diet improved the structure of midgut microvilli compared to shrimp that did not use dietary supplements.

The role of phytochemicals in inhibiting microbial pathogens

The antimicrobial properties of phytochemical compounds vary according to the location of hydroxyl or alkyl groups and the release properties of lipopolysaccharides (Yang *et al.*, 2015). About phenolic terpenoids, the hydroxyl group and the presence of non-localized electrons are important elements for their antimicrobial function (Lambert *et al.*, 2001; Ultee *et al.*, 2002). Hellander *et al.* (1998) also reported that thymol and carvacrol have a high ability to degrade outer membranes. This function is mainly due to their ability to release lipopolysaccharides. Phytochemicals penetrate the bacterial membrane due to their lipophilic properties and disrupt the structure and function of bacteria by expanding the membrane, increasing membrane fluidity and permeability, disrupting membrane proteins, inhibiting respiration, and altering ion transport. (Firmino *et al.*, 2021). These compounds also play a role in reducing infection by detecting and accumulating microbial pathogens through lectin-mediated pathways (Ourth *et al.*, 2007; LaFrentz *et al.*, 2012). Other mechanisms of phytochemicals as antibacterial compounds include inhibition of flagellin synthesis, inhibition of bacterial movement, depletion of ATP reservoirs, elimination

of proton stimulus, inhibition of quorum sensing (QS), and reduction of bacterial biofilm formation (Kachur and Suntres, 2020).

Quorum Sensing (QS) is involved in regulating a wide range of different physiological processes of bacteria, especially in aggravating factors that are important during the pathogenic process (Greenberg, 2003; Vendeville *et al.*, 2005; Xavier and Bassler, 2003; Defoirdt *et al.*, 2013). In this regard, there is evidence for the ability to disrupt QS by various phytochemical compounds including alfalfa seed extract (Vikram *et al.*, 2010), plants (Koh and Tham, 2011), and essential oils (Bjarnsholt *et al.*, 2005; Choo *et al.*, 2006; Zhou *et al.*, 2013). Unfortunately, the use of QS inhibition strategies to control bacterial infections is unknown in aquaculture (Yang *et al.*, 2015). The first report on aquatic animals was made by Manefield *et al.* (1999) on furanone extracted from the Australian microalgae (*Delisea pulchra*). Based on the results of this study, this combination increased the survival of shrimp larvae exposed to pathogens such as *V. campbellii*, *V. harveyi*, and *V. parahaemolyticus* (Defoirdt *et al.*, 2007). Organic compounds such as cinnamaldehyde and its derivatives have also been reported to be effective against *V. harveyi* in Brine shrimp (Niu *et al.*, 2006; Brackman *et al.*, 2008) and *Macrobrachium Rosenbergii* larvae (Pande *et al.*, 2013).

The role of phytogenics in inhibiting microbial pathogens in cold water fish

Studies on the effect of phytogenetics on increasing the resistance of cold-water fish to bacterial pathogens are much less than warm-water fish, and studies have been limited to rainbow trout. In a study by Menanteau-Ledouble *et al.* (2015), the use of a commercial phytogenic food additive (containing oregano, anise, and citrus oils) increased the survival of rainbow trout (*Onchorhynchus mykiss*) exposed to *Aeromonas salmonicida* in three forms of Intraperitoneal injection, immersion in bacterial solution and living with infected fish. According to these results, the highest effect of phytogenics in protecting fish from pathogens was in the form of immersion. Also, according to Gulec *et al.* (2013), fortification of the diet with oils extracted from *Thymus vulgaris* and *Foeniculum vulgare* can increase the resistance to *Yersinia ruckeri* in rainbow trout by increasing the levels of some biochemical parameters and electrolytes.

The role of phytogenics in inhibiting microbial pathogens in warm water fish

Most studies have been performed on the effects of phytogenics on increasing the resistance of tilapia to bacterial diseases. Numerous studies have reported increased resistance of tilapia to *Aeromonas hydrophila* challenge due to the use of phytogenic compounds which include studies by Jayathirtha and Mishra (2004) using *E. officinalis* supplement, El Deen and Razin (2009) using crude garlic extract (*Allium*

sativum), and mugwort (*Artemisia vulgaris*), Gbadamosi *et al.* (2016) using *Moringa oleifera*, and Pachanawan *et al.* (2008) using dry leaf powder and ethanolic extract of *Psidium guajava*. Asimi and Sahu (2013) in the study of feeding with extracts of thyme (*Thymus vulgaris*) rosemary (*Rosmarinus officinalis*) and fenugreek (*Trigonella foenum graecum*) extracts in Mozambique tilapia (*Oreochromis mossambicus*) and Zilberg *et al.* (2010) in the study of feeding with *Rosmarinus officinalis* extracts in Nile tilapia reported increased resistance and decreased fish mortality after infection with *Streptococcus iniae*.

Numerous studies have been performed on catfish, including the effect of using garlic peel (*Allium sativum*) on increasing resistance to *Aeromonas hydrophila* infection in the fingerlings of African catfish (Thanikachalam *et al.*, 2010). Zheng *et al.* (2009) also evaluated the effect of carvacrol and thymol separately, in combination with each other and combination with oregano essential oil in canal catfish. In this study, fish were challenged with this diet for 8 weeks and then with *Aeromonas hydrophila*. After challenge with *A. hydrophila*, the lowest mortality was observed in the group fed with oregano and then the fish fed with thymol and carvacrol. In another study, Peterson *et al.* (2015) found that feeding with a commercial phytogenic feed additive (Digestarom® PEP MGE; containing the essential oils of carvacrol, thymol, anethole, and limonene) increased survival rates in juveniles of

canal catfish (*Ictalurus punctatus*) exposed to *Edwardsiella ictaluri*. Other studies on warm-water fish include Minomol (2005) on the effect of *Emblica officinalis* on reducing the incidence of microbial infection in *Carassius auratus* and the results of Volpatti *et al.* (2013) on resistance of European sea bass fed carvacrol against *Vibrio anguillarum*.

The role of phytochemicals in inhibiting microbial pathogens in shrimp

There are also studies on the protective effect of nutrition with phytochemicals exposed to bacterial pathogens in shrimp. According to Goncalves and Santos (2015), the use of a diet containing the commercial phytochemical compound Digestarom® P.E.P. MGE increased resistance and decreased mortality in shrimp exposed to *Vibrio parahaemolyticus*. Also, according to the report of Yatip *et al.* (2018), the use of fermented Japanese soy extract (Natto) in the diet led to a reduction in mortalities due to *Vibrio harveyi* in white leg shrimp *Penaeus vannamei*.

The role of phytochemicals in inhibiting viral pathogens

The antiviral effects of plant compounds are mainly due to the inhibition of virus transcription, reducing its proliferation in host cells and thus increasing the innate immune response (Syahidah *et al.*, 2015). In this regard, few studies have been performed on the effect of phytochemical food additives in inhibiting viral pathogens in aquatic animals, which is limited to shrimp. The shrimp

farming industry is currently facing several episodic viral diseases. White spot syndrome (WSSV) virus causes 100% mortality in shrimp farms (Selvam *et al.*, 2020). In this regard, according to Yogeewaran *et al.* (2012), the use of *Withania somnifera* (Yogeewaran *et al.*, 2012) and phytochemical supplementation™ Phytocee (Selvam *et al.*, 2020) as a dietary supplement increases the survival of shrimp exposed to White spot disease. However, the results of studies show the antiviral effects of medicinal plants on aquatic viruses and the extensive research capacity to investigate the effects of these plant compounds as food additives in vivo.

For example, in vitro studies include Micol *et al.* (2005) report on the inhibitory effect of olive leaf extract (*Olea europaea*) on Viral hemorrhagic septicemia (VHSV) and the effect of eighteen Thai plant extracts (*Cassia alata*, *Clinacanthus* spp., *Clinacanthus nutans*, *Calophyllum inophyllum*, *Hura crepitans*, *Glinus oppositifolius*, *Momordica charantia*, *Ocimum sanctum* (white), *Ocimum sanctum* (red), *Phyllanthus reticulatus*, *Phyllanthus urinaria*, *Phyllanthus acidus*, *Orchocarpus siamensis*, *Phyllanthus amarus*, *Phyllanthus debilis*, *Psidium guajava*, *Tinospora crispa* and *Tinospora cordifolia*) on CHSE-214 cells were reported in vitro against infectious hematopoietic necrosis virus (IHNV), *Oncorhynchus masou* virus disease (OMV) and Infectious pancreatic necrosis (IPNV) (Direkbusarakom *et al.*, 1996). Also in

the study of Harikrishnan *et al.* (2010) Intraperitoneal injection of aqueous, ethanolic, and methanolic extracts of *Punica granatum* leaves increased resistance to lymphocystis disease virus (LDV) in Olive flounder (*Paralichthys olivaceus*). According to Novriadi and Haw (2015), immersion of tiger grouper (*Epinephelus fuscoguttatus*) in herbal solution © AquaHerb increased fish resistance to iridovirus infection.

The role of phytochemicals in inhibiting parasitic pathogens

Most of the studies on the effect of plant compounds on parasites in aquatic animals have been related to external parasites and the preferred prescription method is immersion due to the lack of proper implementation of prevention methods in aquaculture farms (Reverter *et al.*, 2014). Numerous studies have been done on the inhibitory effects of plant compounds on parasites including *Gyrodactylus turnbulli* (Fridman *et al.*, 2014), *Anacanthorus penilabiatus* (Martins *et al.*, 2002), *Neobenedenia* sp. (Militz *et al.*, 2014), *Ichthyophthirius multifiliis* (Zhang *et al.*, 2013), *Dactylogyrus intermedius* (Wu *et al.*, 2011) and *Gyrodactylus kobayashii* (Zhou *et al.*, 2017). However, limited studies have reported the effects of phytochemical dietary supplements on the prevention and control of parasitic diseases. Existing mechanisms are involved in improving immune function (especially mucosal immunity). Some studies in tilapia have reported a protective effect of phytochemicals against *Trichodina* infection. According to a

study by El Deen and Razin (2009), the use of crude garlic extract (*Allium sativum*) and mugwort (*Artemisia vulgaris*) controlled *Trichodina*'s disease in Nile tilapia. Chitmanat *et al.* (2005) showed that the use of garlic and Indian almond extract (*Terminalia catappa*) reduces *Trichodina* infection in tilapia. Chansue and Tangtrongpiros (2005) also reported that the addition of dried almond leaves (*Terminalia catappa*) could kill monogenic parasites in *Carassius auratus*. According to Fridman *et al.* (2014), the use of the aqueous extract of garlic (*Allium sativum*) in the diet of guinea fowl (*Poecilia reticulata*) infected with *Gyrodactylus turnbulli* and *Dactylogyrus* sp. significantly reduced the mean prevalence and severity of parasites. Militz *et al.* (2013) showed that the use of garlic supplements as prophylaxis leads to a significant reduction in infection of *Neobenedenia* sp. (Monogenea: Capsalidae) in farmed barramundi (*Lates calcarifer*).

The role of phytochemicals in improving the quality of fish carcasses

Carcass composition including fat and protein are factors that have been evaluated in some studies as a result of feeding with phytochemical food additives in aquatic animals. The results of Peterson *et al.* (2014) study on catfish fillets fed with phytochemical feed additive (Digestaron® PEP MGE) showed less fillet fat (31.3 vs. 35.9) and slightly higher fillet protein (62.4 vs. 61.2%). Less fat in seafood is beneficial because it reduces waste, increases efficiency,

reduces bad taste, and reduces storage problems. Also in the study of rainbow trout, the stability of lipids against oxidative damage and the activity of glutathione-based enzymes in the fillets of both thymol-treated fish and carvacrol-treated fish improved (Giannenas *et al.* 2012). The results of these studies show that herbal supplements may have a beneficial effect in reducing fillet fat, increasing protein deposition, and increasing carcass antioxidant activity. However, the results of studies in some cases show the lack of phytoGenic compounds' effect on improving carcass quality. According to the results of Zheng *et al.* (2009), feeding canal catfish with carvacrol and thymol separately, in combination with each other and combination with oregano essential oil did not affect carcasses and fillets among the control and treatment groups.

Prospects for the future

The world's population today is about seven billion and will reach nine billion by 2050. Feeding the world's population by 2050 will be a major challenge. Therefore, aquaculture will replace fishing and most of the required protein will be provided from aquatic sources (Costello *et al.*, 2020). The use of antibiotics and chemicals as stimulants for the growth and treatment of diseases in aquaculture has raised concerns in many countries. These compounds cause environmental pollution and are dangerous to human health. In 2022, several countries and the European Union (EU) will ban the use of

antibiotics on farms. This ban indicates the need to develop alternative therapies (More, 2020). The use of herbal compounds is one of these alternative therapies that have attracted a lot of attention in recent years. The global market for phytoGenic food additives is estimated at \$ 753 million in 2020 and is projected to reach \$ 1.098 million by 2025 (Firmino *et al.*, 2021). Herbal compounds can be effective alternatives, especially for the prevention of diseases caused by bacteria and parasites in aquatic animals. However, studies on this subject have shown varying degrees of effectiveness, and the use of phytoGenic compounds to control parasites through immersion has had limited success. The reason for this can be attributed to factors such as extraction methods, part of the plant that is used as raw material, the concentration of additives, prescription method, and storage conditions before feeding, which greatly affects their performance (Reverter *et al.* 2014). Determining the optimal dose in the use of phytoGenic compounds is also very important because it increases losses if inappropriate doses are used (Ikhwanuddin *et al.*, 2014). Further studies are also needed to determine the quality, bioactive components, and classification of the chemical composition of these phytoGenics to be used to develop standard extraction protocols, determine prescribing methods and determine the optimal dose in fish and crustaceans (Reverter *et al.*, 2014). The fastest way to use phytoGenic compounds is by injection, but this

method is very troublesome and stressful in fish and crustaceans. Therefore, feeding prescription is the most appropriate and available method (Yoshida *et al.*, 1995). One of the valuable initiatives that will help develop the use of these phytogenic compounds in the future is the stabilization of various phytogenic compounds such as volatile essential oils using methods such as matrix-encapsulation. This allows these compounds to remain active in more parts of the gastrointestinal tract (GIT) and their positive effects are not limited to a small part of the GIT (Goncalves and Santos, 2015).

Despite numerous reports of the benefits of using phytogenic compounds in experimental studies, inconsistencies in their effects on host growth and systemic safety and a lack of understanding of how they work in intestinal health and intestinal microbiota have led to limitations in the use of these compounds in aquatic animals. Generally, further studies are needed to introduce and develop sustainable prevention strategies to use phytogenic compounds to enhance host immunity, stress resistance, and the prevention of pathogenesis in aquaculture.

References

- Abdel-Tawwab, M., Adeshina, I., Jenyo-Oni, A., Ajani, E.K. and Emikpe, B.O., 2018.** Growth, physiological, antioxidants, and immune response of African catfish, *Clarias gariepinus* (B.), to dietary clove basil, *Ocimum gratissimum*, leaf extract and its susceptibility to *Listeria monocytogenes* infection. *Fish and Shellfish Immunology*, 78, pp.346-354.
- Abdel-Latif, H.M., Abdel-Tawwab, M., Khafaga, A.F. and Dawood, M.A., 2020.** Dietary oregano essential oil improved the growth performance via enhancing the intestinal morphometry and hepatorenal functions of common carp (*Cyprinus carpio* L.) fingerlings. *Aquaculture*, 526, pp.735432.
- Adel, M., Amiri, A.A., Zorriehzahra, J., Nematollahi, A. and Esteban, M.Á., 2015a.** Effects of dietary peppermint (*Mentha piperita*) on growth performance, chemical body composition and hematological and immune parameters of fry Caspian white fish (*Rutilus frisii kutum*). *Fish and Shellfish Immunology*, 45(2), pp.841-847.
- Adel, M., Safari, R., Pourgholam, R., Zorriehzahra, J. and Esteban, M.Á., 2015b.** Dietary peppermint (*Mentha piperita*) extracts promote growth performance and increase the main humoral immune parameters (both at mucosal and systemic level) of Caspian brown trout (*Salmo trutta caspius* Kessler, 1877). *Fish and Shellfish Immunology*, 47(1), pp.623-629.
- Araruna, M.E., Serafim, C., Alves Júnior, E., Hiruma-Lima, C., Diniz, M. and Batista, L., 2020.** Intestinal Anti-Inflammatory Activity of Terpenes in Experimental

- Models (2010–2020): A Review. *Molecules*, 25(22), p.5430.
- Ardó, L., Yin, G., Xu, P., Váradi, L., Szigeti, G., Jeney, Z. and Jeney, G., 2008.** Chinese herbs (*Astragalus membranaceus* and *Lonicera japonica*) and boron enhance the non-specific immune response of Nile tilapia (*Oreochromis niloticus*) and resistance against *Aeromonas hydrophila*. *Aquaculture*, 275(1-4), pp.26-33.
- Asimi, O.A. and Sahu, N.P., 2013.** Herbs/spices as feed additive in aquaculture. *Scientific Journal of Pure and Applied Sciences*, 2(8), pp.284-292.
- Austin, B., 2006.** The bacterial microflora of fish, revised. *TheScientificWorldJournal*, 6, pp.931-945.
- Bahia, P.K., Rattray, M. and Williams, R.J., 2008.** Dietary flavonoid (–) epicatechin stimulates phosphatidylinositol 3-kinase-dependent anti-oxidant response element activity and up-regulates glutathione in cortical astrocytes. *Journal of Neurochemistry*, 106(5), pp. 2194-2204.
- Baron, M., Davies, S., Alexander, L., Snellgrove, D. and Sloman, K.A., 2008.** The effect of dietary pigments on the coloration and behaviour of flame-red dwarf gourami, *Colisa lalia*. *Animal Behaviour*, 75(3), pp.1041-1051.
- Bello, O.S. and Emikpe, B.O., 2012.** The body weight changes and gut morphometry of *clarias gariepinus* juveniles on feeds supplemented with walnut (*Tetracarpidium conophorum*) leaf and onions (*allium cepa*) bulb residues.
- Bento, M.H.L., Ouwehand, A.C., Tiisonen, K., Lahtinen, S., Nurminen, P., Saarinen, M.T., Schulze, H., Mygind, T. and Fischer, J., 2013.** Essential oils and their use in animal feeds for monogastric animals--Effects on feed quality, gut microbiota, growth performance and food safety: a review. *Veterinari Medicina*, 58(9).
- Bharathi, S., Antony, C., Cbt, R., Arumugam, U., Ahilan, B. and Aanand, S., 2019.** Functional feed additives used in fish feeds. *International Journal of Fisheries and Aquatic Studies*, 7(3), pp.44-52.
- Bhujel, R.C., Poolsawat, L. and Gonçalves, R.A., 2017.** Phytochemical feed additive (PFA) improved survival and growth of Nile tilapia (*Oreochromis niloticus*) fry during Subsequent Nursing. *Journal of Aquaculture in the Tropics*, 32(3/4), pp.413-425.
- Bjarnsholt, T., Jensen, P.Ø., Rasmussen, T.B., Christophersen, L., Calum, H., Hentzer, M., Hougen, H.P., Rygaard, J., Moser, C., Eberl, L. and Høiby, N., 2005.** Garlic blocks quorum sensing and promotes rapid clearing of pulmonary *Pseudomonas aeruginosa* infections. *Microbiology*, 151(12), pp.3873-3880.
- Brackman, G., Defoirdt, T., Miyamoto, C., Bossier, P., Van Calenbergh, S., Nelis, H. and**

- Coenye, T., 2008.** Cinnamaldehyde and cinnamaldehyde derivatives reduce virulence in *Vibrio* spp. by decreasing the DNA-binding activity of the quorum sensing response regulator LuxR. *BMC Microbiology*, 8(1), pp.1-14.
- Ceulemans, S., Coutteau, P. and Robles, R., 2009.** Innovative feed additives improve feed utilization in Nile tilapia. *Global aquaculture advocate*. pp2-9.
- Chansue, N. and Tangtrongpiros, J., 2005.** Effect of dried Indian almond leaf (*Terminalia catappa*) on monogenean parasite of gold fish (*Carassius auratus*). *Wetchasan Sattawaphaet*. 19-28pp.
- Chansue, N., Ponpornpisit, A., Endo, M., Sakai, M. and Yoshida, S., 2000.** Improved immunity of tilapia *Oreochromis niloticus* by C-UP III, a herb medicine. *Fish Pathology*, 35(2), pp.89-90.
- Chitmanat, C., Tongdonmuan, K. and Nunsong, W., 2005.** The use of crude extracts from traditional medicinal plants to eliminate *Trichodina* sp. in tilapia (*Oreochromis niloticus*) fingerlings. *Songklanakarin Journal of Science and Technology*, 27(Suppl 1), pp.359-364.
- Choo, J.H., Rukayadi, Y. and Hwang, J.K., 2006.** Inhibition of bacterial quorum sensing by vanilla extract. *Letters in applied microbiology*, 42(6), pp.637-641.
- Citarasu, T., 2010.** Herbal biomedicines: a new opportunity for aquaculture industry. *Aquaculture International*, 18(3), pp.403-414.
- Costello, C., Cao, L., Gelcich, S., Cisneros-Mata, M.Á., Free, C.M., Froehlich, H.E., Golden, C.D., Ishimura, G., Maier, J., Macadam-Somer, I. and Mangin, T., 2020.** The future of food from the sea. *Nature*, 588(7836), pp.95-100.
- Davies, K. ed., 2009.** Annual plant reviews, plant pigments and their manipulation (Vol. 14). John Wiley and Sons. 120P.
- de Francesco, M., Parisi, G., Médale, F., Lupi, P., Kaushik, S.J. and Poli, B.M., 2004.** Effect of long-term feeding with a plant protein mixture based diet on growth and body/fillet quality traits of large rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 236(1-4), pp.413-429.
- Defoirdt, T., Miyamoto, C.M., Wood, T.K., Meighen, E.A., Sorgeloos, P., Verstraete, W. and Bossier, P., 2007.** The natural furanone (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2 (5H)-furanone disrupts quorum sensing-regulated gene expression in *Vibrio harveyi* by decreasing the DNA-binding activity of the transcriptional regulator protein LuxR. *Environmental Microbiology*, 9(10), pp.2486-2495.
- Defoirdt, T., Brackman, G. and Coenye, T., 2013.** Quorum sensing inhibitors: how strong is the evidence?. *Trends in Microbiology*, 21(12), pp.619-624.
- Direkbusarakom, S., Herunsalee, A., Yoshimizu, M. and Ezura, Y., 1996.** Antiviral activity of several Thai traditional herb extracts against fish

- pathogenic viruses. *Fish Pathology*, 31(4), pp.209-213.
- El Deen, A.I.N. and Razin, A.M., 2009.** Application of some medicinal Plants to eliminate *Trichodina* sp. in tilapia (*Oreochromis niloticus*). *Report and Opinion*, 12(17), p.1.
- El Euony, O.I., Elblehi, S.S., Abdel-Latif, H.M., Abdel-Daim, M.M. and El-Sayed, Y.S., 2020.** Modulatory role of dietary *Thymus vulgaris* essential oil and *Bacillus subtilis* against thiamethoxam-induced hepatorenal damage, oxidative stress, and immunotoxicity in African catfish (*Clarias garipenus*). *Environmental Science and Pollution Research*, 27(18), pp.23108-23128.
- Encarnação, P., 2016.** Functional feed additives in aquaculture feeds. In *Aquafeed formulation* (pp. 217-237). Academic Press.
- Englmaierová, M., Bubancová, I. and Skřivan, M., 2014.** Carotenoids and egg quality. *Acta Fytotechnica et Zootechnica*, 17(2), pp.55-57.
- Enterria, A., Slocum, M., Bengtson, D.A., Karayannakidis, P.D. and Lee, C.M., 2011.** Partial replacement of fish meal with plant protein sources singly and in combination in diets for summer flounder, *Paralichthys dentatus*. *Journal of the World Aquaculture Society*, 42(6), pp.753-765.
- FAO, 2018.** The state of world fisheries and aquaculture 2018—Meeting the sustainable development goals. Rome, Italy: The Food and Agricultural Organization of the United Nations. <http://www.fao.org/3/i9540en/i9540en.pdf>.
- FAO, 2020.** Sustainability in action. *State of World Fisheries and Aquaculture*. Rome, 200.
- Ferreira, P.M., Caldas, D.W., Salaro, A.L., Sartori, S.S., Oliveira, J.M., Cardoso, A.J. and Zuanon, J.A., 2016.** Intestinal and liver morphometry of the Yellow Tail Tetra (*Astyanax altiparanae*) fed with oregano oil. *Anais da Academia Brasileira de Ciências*, 88, pp.911-922.
- Firmino, J.P., Galindo-Villegas, J., Reyes-López, F.E. and Gisbert, E., 2021.** Phytogenic bioactive compounds shape fish mucosal immunity. *Frontiers in Immunology*, 12.
- Franz, C., Baser, K.H.C. and Windisch, W., 2010.** Essential oils and aromatic plants in animal feeding—a European perspective. A review. *Flavour and Fragrance Journal*, 25(5), pp.327-340.
- Fridman, S., Sinai, T. and Zilberg, D., 2014.** Efficacy of garlic based treatments against monogenean parasites infecting the guppy (*Poecilia reticulata* (Peters)). *Veterinary Parasitology*, 203(1-2), pp.51-58.
- Gatlin III, D.M., Barrows, F.T., Brown, P., Dabrowski, K., Gaylord, T.G., Hardy, R.W., Herman, E., Hu, G., Krogdahl, Å., Nelson, R. and Overturf, K., 2007.** Expanding the utilization of sustainable plant products in

- aquafeeds: a review. *Aquaculture Research*, 38(6), pp.551-579.
- Gbadamosi, O.K., Fasakin, A.E. and Adebayo, O.T., 2016.** Hepatoprotective and stress-reducing effects of dietary *Moringa oleifera* extract against *Aeromonas hydrophila* infections and transportation-induced stress in Nile tilapia, *Oreochromis niloticus* (Linnaeus 1757) fingerlings. *International Journal of Environmental and Agriculture Research*, 2, pp.121-128.
- Gessner, D.K., Fiesel, A., Most, E., Dinges, J., Wen, G., Ringseis, R. and Eder, K., 2013.** Supplementation of a grape seed and grape marc meal extract decreases activities of the oxidative stress-responsive transcription factors NF- κ B and Nrf2 in the duodenal mucosa of pigs. *Acta Veterinaria Scandinavica*, 55(1), pp.1-10.
- Gholamhosseini, A., Adel, M., Dawood, M.A. and Banaee, M., 2020.** The potential benefits of *Mentha longifolia* on growth performance and innate immunity parameters in Caspian kutum (*Rutilus frisii kutum*). *Aquaculture Research*, 51(12), pp.5212-5227.
- Giannenas, I., Triantafillou, E., Stavarakakis, S., Margaroni, M., Mavridis, S., Steiner, T. and Karagouni, E., 2012.** Assessment of dietary supplementation with carvacrol or thymol containing feed additives on performance, intestinal microbiota and antioxidant status of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 350, pp.26-32.
- Gómez, G.D. and Balcázar, J.L., 2008.** A review on the interactions between gut microbiota and innate immunity of fish. *FEMS Immunology and Medical Microbiology*, 52(2), pp.145-154.
- Goncalves, R. and Santos, G., 2015.** Phytogenics for better profitability and sustainability. *Growth*, 8, 9 P.
- Greenberg, E.P., 2003.** Bacterial communication and group behavior. *The Journal of clinical investigation*, 112(9), pp.1288-1290.
- Gulec, A.K., Danabas, D., Ural, M., Seker, E., Arslan, A. and Serdar, O., 2013.** Effect of mixed use of thyme and fennel oils on biochemical properties and electrolytes in rainbow trout as a response to *Yersinia ruckeri* infection. *Acta Veterinaria Brno*, 82(3), pp.297-302.
- Hancz, C., Magyary, I., Molnar, T., Sato, S., Horn, P. and Taniguchi, N., 2003.** Evaluation of color intensity enhanced by paprika as feed additive in goldfish and koi carp using computer-assisted image analysis. *Fisheries Science*, 69(6), pp.1158-1161.
- Hardy, R.W., 2010.** Utilization of plant proteins in fish diets: effects of global demand and supplies of fishmeal. *Aquaculture Research*, 41(5), pp.770-776.
- Harikrishnan, R., Heo, J., Balasundaram, C., Kim, M.C., Kim, J.S., Han, Y.J. and Heo, M.S., 2010.** Effect of *Punica granatum* solvent extracts on immune system

- and disease resistance in *Paralichthys olivaceus* against lymphocystis disease virus (LDV). *Fish and Shellfish Immunology*, 29(4), pp.668-673.
- He, W., Rahimnejad, S., Wang, L., Song, K., Lu, K. and Zhang, C., 2017.** Effects of organic acids and essential oils blend on growth, gut microbiota, immune response and disease resistance of Pacific white shrimp (*Litopenaeus vannamei*) against *Vibrio parahaemolyticus*. *Fish and Shellfish Immunology*, 70, pp.164-173.
- Helander, I.M., Alakomi, H.L., Latva-Kala, K., Mattila-Sandholm, T., Pol, I., Smid, E.J., Gorris, L.G. and von Wright, A., 1998.** Characterization of the action of selected essential oil components on Gram-negative bacteria. *Journal of Agricultural and Food Chemistry*, 46(9), pp.3590-3595.
- Heluy, G.M., Ramos, L.R.V., Pedrosa, V.F., Sarturi, C., Figueiredo, P.G.P., Vidal, L.G.P., Franca, I.D.F. and Pereira, M.M., 2020.** Oregano (*Origanum vulgare*) essential oil as an additive in diets for Nile tilapia (*Oreochromis niloticus*) fingerlings reared in salinized water. *Aquaculture Research*, 51(8), pp.3237-3243.
- Hernández, A., García, B.G., Caballero, M.J. and Hernández, M.D., 2015.** Preliminary insights into the incorporation of rosemary extract (*Rosmarinus officinalis* L.) in fish feed: influence on performance and physiology of gilthead seabream (*Sparus aurata*). *Fish Physiology and Biochemistry*, 41(4), pp.1065-1074.
- Hernández, A., García, B., Caballero, M.J. and Hernández, M.D., 2016.** The inclusion of thyme essential oil in the feed of gilthead seabream (*Sparus aurata*) promotes changes in the frequency of lymphocyte aggregates in gut-associated lymphoid tissue. *Aquaculture Research*, 47(10), pp.3341-3345.
- Heydari, M., Firouzbakhsh, F. and Paknejad, H., 2020.** Effects of *Mentha longifolia* extract on some blood and immune parameters, and disease resistance against yersiniosis in rainbow trout. *Aquaculture*, 515, p.734586.
- Heyman, H.M., Senejoux, F., Seibert, I., Klimkait, T., Maharaj, V.J. and Meyer, J.J.M., 2015.** Identification of anti-HIV active dicaffeoylquinic- and tricaffeoylquinic acids in *Helichrysum populifolium* by NMR-based metabolomic guided fractionation. *Fitoterapia*, 103, pp.155-164.
- Hong, J.C., Steiner, T., Aufy, A. and Lien, T.F., 2012.** Effects of supplemental essential oil on growth performance, lipid metabolites and immunity, intestinal characteristics, microbiota and carcass traits in broilers. *Livestock Science*, 144(3), pp.253-262.
- Hongbai, L., Ying, Z., Yuhui, Y.A.N.G., Tongyan, L. and Jidan, Y., 2004.** Effects of five Chinese herb medicines as additive in feed on the growth and intestinal microflora in

- common carp (*Cyprinus carpio*). *Journal of Dalian Fisheries College*, 19(1), pp.16-20.
- Hoseini, S.M. and Yousefi, M., 2019.** Beneficial effects of thyme (*Thymus vulgaris*) extract on oxytetracycline-induced stress response, immunosuppression, oxidative stress and enzymatic changes in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Nutrition*, 25(2), pp.298-309.
- Hoseinifar, S.H., Zou, H.K., Miandare, H.K., Van Doan, H., Romano, N. and Dadar, M., 2017.** Enrichment of common carp (*Cyprinus carpio*) diet with medlar (*Mespilus germanica*) leaf extract: Effects on skin mucosal immunity and growth performance. *Fish and Shellfish Immunology*, 67, pp.346-352.
- Hwang, Y.P., Choi, J.H., Yun, H.J., Han, E.H., Kim, H.G., Kim, J.Y., Park, B.H., Khanal, T., Choi, J.M., Chung, Y.C. and Jeong, H.G., 2011.** Anthocyanins from purple sweet potato attenuate dimethylnitrosamine-induced liver injury in rats by inducing Nrf2-mediated antioxidant enzymes and reducing COX-2 and iNOS expression. *Food and Chemical Toxicology*, 49(1), pp.93-99.
- Ikhwanuddin, M., Moh, J.H., Hidayah, M., Noor-Hidayati, A.B., Aina-Lyana, N.M. and Juneta, A.S., 2014.** Effect of Indian almond, *Terminalia catappa* leaves water extract on the survival rate and growth performance of black tiger shrimp, *Penaeus monodon* post larvae. *Aquaculture, Aquarium, Conservation and Legislation*, 7(2), pp.85-93.
- Immanuel, G., Vincybai, V.C., Sivaram, V., Palavesam, A. and Marian, M.P., 2004.** Effect of butanolic extracts from terrestrial herbs and seaweeds on the survival, growth and pathogen (*Vibrio parahaemolyticus*) load on shrimp *Penaeus indicus* juveniles. *Aquaculture*, 236(1-4), pp.53-65.
- Jayathirtha, M.G. and Mishra, S.H., 2004.** Preliminary immunomodulatory activities of methanol extracts of *Eclipta alba* and *Centella asiatica*. *Phytomedicine*, 11(4), pp.361-365.
- Kachur, K. and Suntres, Z., 2020.** The antibacterial properties of phenolic isomers, carvacrol and thymol. *Critical Reviews in Food Science and Nutrition*, 60(18), pp.3042-3053.
- Karásková, K., Suchý, P. and Straková, E., 2015.** Current use of phytogenic feed additives in animal nutrition: a review. *Czech Journal of Animal Science*, 60(12), pp.521-30.
- Kesselring, J., Gruber, C., Standen, B. and Wein, S., 2021.** Effect of a phytogenic feed additive on the growth performance and immunity of Pacific white leg shrimp, *Litopenaeus vannamei*, fed a low fishmeal diet. *Journal of the World Aquaculture Society*, 52(2), pp.303-315.
- Khafaga, A.F., Naiel, M.A., Dawood, M.A. and Abdel-Latif, H.M., 2020.**

- Dietary *Origanum vulgare* essential oil attenuates cypermethrin-induced biochemical changes, oxidative stress, histopathological alterations, apoptosis, and reduces DNA damage in Common carp (*Cyprinus carpio*). *Aquatic Toxicology*, 228, p.105624.
- Koh, K.H. and Tham, F.Y., 2011.** Screening of traditional Chinese medicinal plants for quorum-sensing inhibitors activity. *Journal of Microbiology, Immunology and Infection*, 44(2), pp.144-148.
- Korzekwa, M., Wilke, T. and Brenner, S., 2016.** Positive effects of a flavonoid-rich phytochemical product on growth, survival, and bacterial infection of Pacific White Shrimp. In 15. *BOKU-Symposium TIERERNÄHRUNG*.
- Kroismayr, A., Steiner, T. and Zhang, C., 2006.** Influence of a phytochemical feed additive on performance of weaner piglets. In *Journal of Animal Science* (Vol. 84, pp. 270-270). 1111 NORTH DUNLAP AVE, SAVOY, IL 61874 USA: AMER SOC ANIMAL SCIENCE.
- Kurian, A., Van Doan, H., Tapingkae, W. and Elumalai, P., 2020.** Modulation of mucosal parameters, innate immunity, growth and resistance against *Streptococcus agalactiae* by enrichment of Nile tilapia (*Oreochromis niloticus*) diet with *Leucas aspera*. *Fish and Shellfish Immunology*, 97, pp.165-172.
- LaFrentz, B.R., Booth, N.J., Ourth, D.D. and Peterson, B.C., 2012.** Spleen size and plasma levels of mannose binding lectin in channel catfish families challenged with *Flavobacterium columnare* and *Edwardsiella ictaluri*. *Journal of Aquatic Animal Health*, 24, pp.141-147.
- Laltlanmawia, C., Saha, R.K., Saha, H. and Biswas, P., 2019.** Ameliorating effects of dietary mixture of *Withania somnifera* root extract and vitamin C in *Labeo rohita* against low pH and waterborne iron stresses. *Fish and Shellfish Immunology*, 88, pp.170-178.
- Lambert, R.J.W., Skandamis, P.N., Coote, P.J. and Nychas, G.J., 2001.** A study of the minimum inhibitory concentration and mode of action of oregano essential oil, thymol and carvacrol. *Journal of Applied Microbiology*, 91(3), pp.453-462.
- Lazado, C.C. and Caipang, C.M.A., 2014.** Mucosal immunity and probiotics in fish. *Fish and shellfish immunology*, 39(1), pp.78-89.
- Lee, Y., Shin, D.H., Kim, J.H., Hong, S., Choi, D., Kim, Y.J., Kwak, M.K. and Jung, Y., 2010.** Caffeic acid phenethyl ester-mediated Nrf2 activation and I κ B kinase inhibition are involved in NF κ B inhibitory effect: structural analysis for NF κ B inhibition. *European Journal of Pharmacology*, 643(1), pp.21-28.
- Liu, H.W., Tong, J.M. and Zhou, D.W., 2011.** Utilization of Chinese herbal feed additives in animal production. *Agricultural Sciences in China*, 10(8), pp.1262-1272.

- Liu, S.D., Song, M.H., Yun, W., Lee, J.H., Kim, H.B. and Cho, J.H., 2019.** Effect of carvacrol essential oils on immune response and inflammation-related genes expression in broilers challenged by lipopolysaccharide. *Poultry Science*, 98(5), pp.2026-2033.
- Llana-Ruiz-Cabello, M., Gutiérrez-Praena, D., Puerto, M., Pichardo, S., Jos, Á. and Cameán, A.M., 2015.** In vitro pro-oxidant/antioxidant role of carvacrol, thymol and their mixture in the intestinal Caco-2 cell line. *Toxicology in Vitro*, 29(4), pp.647-656.
- Luo, R., 1997.** Induction of immunity substance in *Penaeus chinensis* by Chinese herbal medicine. *Oceanologia et Limnologia Sinica*, 28, pp.577-585.
- Manefield, M., de Nys, R., Naresh, K., Roger, R., Givskov, M., Peter, S. and Kjelleberg, S., 1999.** Evidence that halogenated furanones from *Delisea pulchra* inhibit acylated homoserine lactone (AHL)-mediated gene expression by displacing the AHL signal from its receptor protein. *Microbiology*, 145(2), pp.283-291.
- Martins, M.L., Moraes, F.R., Miyazaki, D.M.Y., Brum, C.D., Onaka, E.M., Fenerick, J.J. and Bozzo, F.R., 2002.** Alternative treatment for *Anacanthorus penilabiatus* (Monogenea: Dactylogyridae) infection in cultivated pacu, *Piaractus mesopotamicus* (Osteichthyes: Characidae) in Brazil and its haematological effects. *Parasite*, 9(2), pp.175-180.
- McDevitt, R.M., Brooker, J.D., Acamovic, T. and Sparks, N.H.C., 2006.** Necrotic enteritis; a continuing challenge for the poultry industry. *World's Poultry Science Journal*, 62(2), pp.221-247.
- Menanteau-Ledouble, S., Krauss, I., Santos, G., Fibi, S., Weber, B. and El-Matbouli, M., 2015.** Effect of a phytogenic feed additive on the susceptibility of *Onchorhynchus mykiss* to *Aeromonas salmonicida*. *Diseases of Aquatic Organisms*, 115(1), pp.57-66.
- Micol, V., Caturla, N., Pérez-Fons, L., Más, V., Pérez, L. and Estepa, A., 2005.** The olive leaf extract exhibits antiviral activity against viral haemorrhagic septicemia rhabdovirus (VHSV). *Antiviral Research*, 66(2-3), pp.129-136.
- Militz, T.A., Southgate, P.C., Carton, A.G. and Hutson, K.S., 2014.** Efficacy of garlic (*Allium sativum*) extract applied as a therapeutic immersion treatment for *Neobenedenia* sp. management in aquaculture. *Journal of Fish Diseases*, 37(5), pp.451-461.
- Militz, T.A., Southgate, P.C., Carton, A.G. and Hutson, K.S., 2013.** Dietary supplementation of garlic (*Allium sativum*) to prevent monogenean infection in aquaculture. *Aquaculture*, 408, pp.95-99.
- Minomol, M., 2005.** Culture of Gold fish *Carassius auratus* using medicinal plants having immunostimulant characteristics

- (Doctoral dissertation, M. Phil Dissertation, MS University, India).
- Mirghaed, A.T., Hoseini, S.M., Hoseinifar, S.H. and Van Doan, H., 2020.** Effects of dietary thyme (*Zataria multiflora*) extract on antioxidant and immunological responses and immune-related gene expression of rainbow trout (*Oncorhynchus mykiss*) juveniles. *Fish and Shellfish Immunology*, 106, pp.502-509.
- Mohammadi, G., Rafiee, G., El Basuini, M.F., Van Doan, H., Ahmed, H.A., Dawood, M.A. and Abdel-Latif, H.M., 2020.** Oregano (*Origanum vulgare*), St John's-wort (*Hypericum perforatum*), and lemon balm (*Melissa officinalis*) extracts improved the growth rate, antioxidative, and immunological responses in Nile tilapia (*Oreochromis niloticus*) infected with *Aeromonas hydrophila*. *Aquaculture Reports*, 18, pp.100445.
- Montero, D., Robaina, L., Caballero, M.J., Gines, R. and Izquierdo, M.S., 2005.** Growth, feed utilization and flesh quality of European sea bass (*Dicentrarchus labrax*) fed diets containing vegetable oils: A time-course study on the effect of a re-feeding period with a 100% fish oil diet. *Aquaculture*, 248(1-4), pp.121-134.
- More, S.J., 2020.** European perspectives on efforts to reduce antimicrobial usage in food animal production. *Irish Veterinary Journal*, 73(1), pp.1-12.
- Mousavi, E., Mohammadi, H., Mousavi, S.M. and Ghatrami, E.R., 2016.** Effects of inulin, savory and onion powders in diet of juveniles carp *Cyprinus carpio* (Linnaeus 1758) on gut micro flora, immune response and blood biochemical parameters. *Turkish Journal of Fisheries and Aquatic Sciences*, 16(4), pp.831-838.
- Nahak, G. and Sahu, R., 2014.** Immunostimulatory effects of *Ocimum sanctum* Linn. leaf extracts in *Clarias batrachus* Linn. *Asian Journal of Pharmaceutical and Clinical Research*, 7(3), pp.157-63.
- Navarrete, P., Toledo, I., Mardones, P., Opazo, R., Espejo, R. and Romero, J., 2010.** Effect of *Thymus vulgaris* essential oil on intestinal bacterial microbiota of rainbow trout, *Oncorhynchus mykiss* (Walbaum) and bacterial isolates. *Aquaculture Research*, 41(10), pp.e667-e678.
- Niu, C., Afre, S. and Gilbert, E.S., 2006.** Subinhibitory concentrations of cinnamaldehyde interfere with quorum sensing. *Letters in Applied Microbiology*, 43(5), pp.489-494.
- Novriadi, R. and Haw, K.B., 2015.** Preliminary study: The use of herbal extracts against iridovirus in tiger grouper *Epinephelus Fuscoguttatus* culture. *Journal of Medicinal Plants*, 3(4), pp.115-120.
- Olsen, R.L. and Hasan, M.R., 2012.** A limited supply of fishmeal: Impact on future increases in global aquaculture production. *Trends in Food Science and Technology*, 27(2), pp.120-128.

- Omoregie, E., Igoche, L., Ojobe, T.O., Absalom, K.V. and Onusiriuka, B.C., 2009.** Effect of varying levels of sweet potato (*Ipomea Batatas*) peels on growth, feed utilization and some biochemical responses of the cichlid (*Oreochromis Niloticus*). *African Journal of Food, Agriculture, Nutrition and Development*, 9(2), pp.700-712.
- Ourth, D.D., Narra, M.B. and Simco, B.A., 2007.** Comparative study of mannose-binding C-type lectin isolated from channel catfish (*Ictalurus punctatus*) and blue catfish (*Ictalurus furcatus*). *Fish and Shellfish Immunology*, 23(6), pp.1152-1160.
- Pachanawan, A., Phumkhachorn, P. and Rattanachaikunsopon, P., 2008.** Potential of Psidium guajava supplemented fish diets in controlling *Aeromonas hydrophila* infection in tilapia (*Oreochromis niloticus*). *Journal of Bioscience and Bioengineering*, 106(5), pp.419-424.
- Paknejad, H., Shekarabi, S.P.H., Mehrgan, M.S., Hajimoradloo, A., Khorshidi, Z. and Rastegari, S., 2020.** Dietary peppermint (*Mentha piperita*) powder affects growth performance, hematological indices, skin mucosal immune parameters, and expression of growth and stress-related genes in Caspian roach (*Rutilus caspicus*). *Fish Physiology and Biochemistry*, 46(5), pp.1883-1895.
- Pande, G.S.J., Scheie, A.A., Benneche, T., Wille, M., Sorgeloos, P., Bossier, P. and Defoirdt, T., 2013.** Quorum sensing-disrupting compounds protect larvae of the giant freshwater prawn *Macrobrachium rosenbergii* from *Vibrio harveyi* infection. *Aquaculture*, 406, pp.121-124.
- Pandey, A.K., Kumar, P. and Saxena, M.J., 2019.** Feed additives in animal health. In *Nutraceuticals in Veterinary Medicine* (pp. 345-362). Springer, Cham.
- Peñaflorida, V.D., 1995.** Growth and survival of juvenile tiger shrimp fed diet where fish meal is partially replaced with papaya (*Carica papaya* L.) or camote (*Ipomea batatas* Lam.) leaf meal. *The Israeli Journal of Aquaculture-Bamidgeh*, 47(1), pp.25-33.
- Peres, H. and Oliva-Teles, A., 2006.** Effect of the dietary essential to non-essential amino acid ratio on growth, feed utilization and nitrogen metabolism of European sea bass (*Dicentrarchus labrax*). *Aquaculture*, 256(1-4), pp.395-402.
- Peterson, B.C., Bosworth, B.G., Li, M.H., Beltran, R. and Santos, G.A., 2014.** Assessment of a phytogenic feed additive (Digestaron PEP MGE) on growth performance, processing yield, fillet composition, and survival of channel catfish. *Journal of the World Aquaculture Society*, 45(2), pp.206-212.
- Peterson, B.C., Peatman, E.R.I.C., Ourth, D.D. and Waldbieser, G.C., 2015.** Effects of a phytogenic feed additive on growth performance,

- susceptibility of channel catfish to *Edwardsiella ictaluri* and levels of mannose binding lectin. *Fish and Shellfish Immunology*, 44(1), pp.21-25.
- Ran, C., Hu, J., Liu, W., Liu, Z., He, S., Dan, B.C.T., Diem, N.N., Ooi, E.L. and Zhou, Z., 2016.** Thymol and carvacrol affect hybrid tilapia through the combination of direct stimulation and an intestinal microbiota-mediated effect: insights from a germ-free zebrafish model. *The Journal of Nutrition*, 146(5), pp.1132-1140.
- Rao, Y.V., Das, B.K., Jyotirmayee, P. and Chakrabarti, R., 2006.** Effect of *Achyranthes aspera* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Fish and Shellfish Immunology*, 20(3), pp.263-273.
- Rashidian, G., Boldaji, J.T., Rainis, S., Prokić, M.D. and Faggio, C., 2021.** Oregano (*Origanum vulgare*) Extract Enhances Zebrafish (*Danio rerio*) Growth Performance, Serum and Mucus Innate Immune Responses and Resistance against *Aeromonas hydrophila* Challenge. *Animals*, 11(2), p.299.
- Reveco, F.E., Øverland, M., Romarheim, O.H. and Mydland, L.T., 2014.** Intestinal bacterial community structure differs between healthy and inflamed intestines in Atlantic salmon (*Salmo salar* L.). *Aquaculture*, 420, pp.262-269.
- Reverter, M., Bontemps, N., Lecchini, D., Banaigs, B. and Sasal, P., 2014.** Use of plant extracts in fish aquaculture as an alternative to chemotherapy: current status and future perspectives. *Aquaculture*, 433, pp.50-61.
- Reyes-Cerpa, S., Vallejos-Vidal, E., Gonzalez-Bown, M.J., Morales-Reyes, J., Pérez-Stuardo, D., Vargas, D., Imarai, M., Cifuentes, V., Spencer, E., Sandino, A.M. and Reyes-López, F.E., 2018.** Effect of yeast (*Xanthophyllomyces dendrorhous*) and plant (Saint John's wort, lemon balm, and rosemary) extract based functional diets on antioxidant and immune status of Atlantic salmon (*Salmo salar*) subjected to crowding stress. *Fish and Shellfish Immunology*, 74, pp.250-259.
- Ringø, E., Sperstad, S., Kraugerud, O.F. and Krogdahl, Å., 2008.** Use of 16S rRNA gene sequencing analysis to characterize culturable intestinal bacteria in Atlantic salmon (*Salmo salar*) fed diets with cellulose or non-starch polysaccharides from soy. *Aquaculture Research*, 39(10), pp.1087-1100.
- Sahu, S., Das, B.K., Mishra, B.K., Pradhan, J. and Sarangi, N., 2007.** Effect of *Allium sativum* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Journal of Applied Ichthyology*, 23(1), pp.80-86.
- Selvam, R., Marimuthu, S. and D'Souza, P., 2020.** Impact of phytocee™, a phytogenic feed additive on survivability of *Litopenaeus vannamei* shrimps under

- WSSV and salinity stress challenges. *International Journal of Fisheries and Aquatic Studies*, 8, pp.261-265.
- Sharma, A., Deo, A.D., Riteshkumar, S.T., Chanu, T.I. and Das, A., 2010.** Effect of *Withania somnifera* (L. Dunal) root as a feed additive on immunological parameters and disease resistance to *Aeromonas hydrophila* in *Labeo rohita* (Hamilton) fingerlings. *Fish and Shellfish Immunology*, 29(3), pp.508-512.
- Shehzad, A., Ha, T., Subhan, F. and Lee, Y.S., 2011.** New mechanisms and the anti-inflammatory role of curcumin in obesity and obesity-related metabolic diseases. *European Journal of Nutrition*, 50(3), pp.151-161.
- Silva, F.C.D.P., Nicoli, J.R., Zambonino-Infante, J.L., Kaushik, S. and Gatesoupe, F.J., 2011.** Influence of the diet on the microbial diversity of faecal and gastrointestinal contents in gilthead sea bream (*Sparus aurata*) and intestinal contents in goldfish (*Carassius auratus*). *FEMS Microbiology Ecology*, 78(2), pp.285-296.
- Slater, M., D'Abramo, L. and Engle, C.R., 2018.** Aquaculture research priorities for the next decade: a global perspective.
- Solà-Oriol, D., Roura, E. and Torrallardona, D., 2011.** Feed preference in pigs: Effect of selected protein, fat, and fiber sources at different inclusion rates. *Journal of Animal Science*, 89(10), pp.3219-3227.
- Somensi, N., Rabelo, T.K., Guimaraes, A.G., Quintans-Junior, L.J., de Souza Araújo, A.A., Moreira, J.C.F. and Gelain, D.P., 2019.** Carvacrol suppresses LPS-induced pro-inflammatory activation in RAW 264.7 macrophages through ERK1/2 and NF-kB pathway. *International Immunopharmacology*, 75, p.105743.
- Steiner, T. and Syed, B., 2015.** Phytogenic feed additives in animal nutrition. In *Medicinal and aromatic plants of the world* (pp. 403-423). Springer, Dordrecht.
- Sudaryono, A., And, E.T. and Evans, L.H., 1999.** Replacement of soybean meal by lupin meal in practical diets for juvenile *Penaeus monodon*. *Journal of the World Aquaculture Society*, 30(1), pp.46-57.
- Sutili, F.J., Velasquez, A., Pinheiro, C.G., Heinzmann, B.M., Gatlin III, D.M. and Baldisserotto, B., 2016.** Evaluation of *Ocimum americanum* essential oil as an additive in red drum (*Sciaenops ocellatus*) diets. *Fish and Shellfish Immunology*, 56, pp.155-161.
- Sutili, F.J., Gatlin III, D.M., Heinzmann, B.M. and Baldisserotto, B., 2018.** Plant essential oils as fish diet additives: benefits on fish health and stability in feed. *Reviews in Aquaculture*, 10(3), pp.716-726.

- Syahidah, A., Saad, C.R., Daud, H.M. and Abdelhadi, Y.M., 2015.** Status and potential of herbal applications in aquaculture: A review.
- Teimouri, M., Amirkolaie, A.K. and Yeganeh, S., 2013.** Effect of *Spirulina platensis* meal as a feed supplement on growth performance and pigmentation of rainbow trout (*Oncorhynchus mykiss*). *World Journal of Fish and Marine Sciences*, 5(2), pp.194-202.
- Thanikachalam, K., Kasi, M. and Rathinam, X., 2010.** Effect of garlic peel on growth, hematological parameters and disease resistance against *Aeromonas hydrophila* in African catfish *Clarias gariepinus* (Bloch) fingerlings. *Asian Pacific Journal of Tropical Medicine*, 3(8), pp.614-618.
- Ultee, A., Bennik, M.H.J. and Moezelaar, R.J.A.E.M., 2002.** The phenolic hydroxyl group of carvacrol is essential for action against the food-borne pathogen *Bacillus cereus*. *Applied and Environmental Microbiology*, 68(4), pp.1561-1568.
- Valladao, G.M., Gallani, S.U., Pala, G., Jesus, R.B., Kotzent, S., Costa, J.C., Silva, T.F. and Pilarski, F., 2017.** Practical diets with essential oils of plants activate the complement system and alter the intestinal morphology of Nile tilapia. *Aquaculture Research*, 48(11), pp.5640-5649.
- Valladão, G.M.R., Gallani, S.U., Kotzent, S., Assane, I.M. and Pilarski, F., 2019.** Effects of dietary thyme essential oil on hemato-immunological indices, intestinal morphology, and microbiota of Nile tilapia. *Aquaculture International*, 27(2), pp.399-411.
- Van de Braak, C.B.T., Botterblom, M.H.A., Taverne, N.V., Van Muiswinkel, W.B., Rombout, J.H.W.M. and Van der Knaap, W.P.W., 2002.** The roles of haemocytes and the lymphoid organ in the clearance of injected *Vibrio* bacteria in *Penaeus monodon* shrimp. *Fish and Shellfish Immunology*, 13(4), pp.293-309.
- Vazirzadeh, A., Jalali, S. and Farhadi, A., 2019.** Antibacterial activity of *Oliveria decumbens* against *Streptococcus iniae* in Nile tilapia (*Oreochromis niloticus*) and its effects on serum and mucosal immunity and antioxidant status. *Fish and Shellfish Immunology*, 94, pp.407-416.
- Veit, M., Jungbauer, L., Wendler, K.R. and Zentner, E., 2011.** Effects of phytochemical feed additives containing *Quillaja saponaria* on ammonia in fattening pigs. In *Animal hygiene and sustainable livestock production. Proceedings of the XVth International Congress of the International Society for Animal Hygiene, Vienna, Austria, 3-7 July 2011, Volume 3* (pp. 1255-1257). Tribuna EU.
- Vendeville, A., Winzer, K., Heurlier, K., Tang, C.M. and Hardie, K.R., 2005.** Making sense of metabolism: autoinducer-2, LuxS and pathogenic bacteria. *Nature Reviews Microbiology*, 3(5), pp.383-396.

- Venkatramalingam, K., Christopher, J.G. and Citarasu, T., 2007.** *Zingiber officinalis* an herbal appetizer in the tiger shrimp *Penaeus monodon* (Fabricius) larviculture. *Aquaculture Nutrition*, 13(6), pp.439-443.
- Vickers, N.J., 2017.** Animal communication: when i'm calling you, will you answer too?. *Current Biology*, 27(14), pp.R713-R715.
- Vikram, A., Jayaprakasha, G.K., Jesudhasan, P.R., Pillai, S.D. and Patil, B.S., 2010.** Suppression of bacterial cell-cell signalling, biofilm formation and type III secretion system by citrus flavonoids. *Journal of Applied Microbiology*, 109(2), pp.515-527.
- Volpatti, D., Chiara, B., Francesca, T. and Marco, G., 2013.** Growth parameters, innate immune response and resistance to *Listonella (Vibrio) anguillarum* of *Dicentrarchus labrax* fed carvacrol supplemented diets. *Aquaculture Research*, 45(1), pp.31-44.
- Waagbø, R. and Remø, S.C., 2020.** Functional diets in fish health management. In *Aquaculture Health Management* (pp. 187-234). Academic Press.
- Wenk, C., 2003.** Herbs and botanicals as feed additives in monogastric animals. *Asian-Australasian Journal of Animal Sciences*, 16(2), pp.282-289.
- Wondrak, G.T., Villeneuve, N.F., Lamore, S.D., Bause, A.S., Jiang, T. and Zhang, D.D., 2010.** The cinnamon-derived dietary factor cinnamic aldehyde activates the Nrf2-dependent antioxidant response in human epithelial colon cells. *Molecules*, 15(5), pp.3338-3355.
- Wong, S. and Rawls, J.F., 2012.** Intestinal microbiota composition in fishes is influenced by host ecology and environment, 3100-3102.
- Wu, Z.F., Zhu, B., Wang, Y., Lu, C. and Wang, G.X., 2011.** In vivo evaluation of anthelmintic potential of medicinal plant extracts against *Dactylogyrus intermedius* (Monogenea) in goldfish (*Carassius auratus*). *Parasitology Research*, 108(6), pp.1557-1563.
- Wu, H., Jiang, K., Yin, N., Ma, X., Zhao, G., Qiu, C. and Deng, G., 2017.** Thymol mitigates lipopolysaccharide-induced endometritis by regulating the TLR4- and ROS-mediated NF-κB signaling pathways. *Oncotarget*, 8(12), p.20042.
- Xavier, K.B. and Bassler, B.L., 2003.** LuxS quorum sensing: more than just a numbers game. *Current Opinion in Microbiology*, 6(2), pp.191-197.
- Yang, C., Chowdhury, M.A., Huo, Y. and Gong, J., 2015.** Phytogetic compounds as alternatives to in-feed antibiotics: potentials and challenges in application. *Pathogens*, 4(1), pp.137-156.
- Yatip, P., Teja, D.N.C., Flegel, T.W. and Soowannayan, C., 2018.** Extract from the fermented soybean product Natto inhibits *Vibrio* biofilm formation and reduces shrimp mortality from *Vibrio harveyi*

- infection. *Fish and Shellfish Immunology*, 72, pp.348-355.
- Yegani, M. and Korver, D.R., 2008.** Factors affecting intestinal health in poultry. *Poultry Science*, 87(10), pp.2052-2063.
- Yeh, R.Y., Shiu, Y.L., Shei, S.C., Cheng, S.C., Huang, S.Y., Lin, J.C. and Liu, C.H., 2009.** Evaluation of the antibacterial activity of leaf and twig extracts of stout camphor tree, *Cinnamomum kanehirae*, and the effects on immunity and disease resistance of white shrimp, *Litopenaeus vannamei*. *Fish and Shellfish Immunology*, 27(1), pp.26-32.
- Yin, G., Jeney, G., Racz, T., Xu, P., Jun, X. and Jeney, Z., 2006.** Effect of two Chinese herbs (*Astragalus radix* and *Scutellaria radix*) on non-specific immune response of tilapia, *Oreochromis niloticus*. *Aquaculture*, 253(1-4), pp.39-47.
- Yogeeswaran, A., Velmurugan, S., Punitha, S.M.J., Babu, M.M., Selvaraj, T., Kumaran, T. and Citarasu, T., 2012.** Protection of *Penaeus monodon* against white spot syndrome virus by inactivated vaccine with herbal immunostimulants. *Fish and Shellfish Immunology*, 32(6), pp.1058-1067.
- Yoshida, T., Kruger, R. and Inglis, V., 1995.** Augmentation of non-specific protection in African catfish, *Clarias gariepinus* (Burchell), by the long-term oral administration of immunostimulants. *Journal of Fish Diseases*, 18(2), pp.195-198.
- Yousefi, M., Ghafarifarsani, H., Hoseinifar, S.H., Rashidian, G. and Van Doan, H., 2021.** Effects of dietary marjoram, *Origanum majorana* extract on growth performance, hematological, antioxidant, humoral and mucosal immune responses, and resistance of common carp, *Cyprinus carpio* against *Aeromonas hydrophila*. *Fish and Shellfish Immunology*, 108, pp.127-133.
- Yu, M.C., Li, Z.J., Lin, H.Z., Wen, G.L. and Ma, S., 2008.** Effects of dietary Bacillus and medicinal herbs on the growth, digestive enzyme activity, and serum biochemical parameters of the shrimp *Litopenaeus vannamei*. *Aquaculture International*, 16(5), pp.471-480.
- Yu, Y., Wang, Q., Huang, Z., Ding, L. and Xu, Z., 2020.** Immunoglobulins, mucosal immunity and vaccination in teleost fish. *Frontiers in Immunology*, 11, p.2597.
- Yuan, C., Pan, X., Gong, Y., Xia, A., Wu, G., Tang, J. and Han, X., 2008.** Effects of Astragalus polysaccharides (APS) on the expression of immune response genes in head kidney, gill and spleen of the common carp, *Cyprinus carpio* L. *International Immunopharmacology*, 8(1), pp.51-58.
- Zarkasi, K.Z., Taylor, R.S., Glencross, B.D., Abell, G.C., Tamplin, M.L. and Bowman, J.P., 2017.** In vitro characteristics of an Atlantic salmon (*Salmo salar* L.) hind

- gut microbial community in relation to different dietary treatments. *Research in Microbiology*, 168(8), pp.751-759.
- Zhang, Q., Xu, D.H. and Klesius, P.H., 2013.** Evaluation of an antiparasitic compound extracted from *Galla chinensis* against fish parasite *Ichthyophthirius multifiliis*. *Veterinary Parasitology*, 198(1-2), pp.45-53.
- Zhang, R., Wang, X.W., Liu, L.L., Cao, Y.C. and Zhu, H., 2020.** Dietary oregano essential oil improved the immune response, activity of digestive enzymes, and intestinal microbiota of the koi carp, *Cyprinus carpio*. *Aquaculture*, 518, pp.734781.
- Zheng, Z.L., Tan, J.Y., Liu, H.Y., Zhou, X.H., Xiang, X. and Wang, K.Y., 2009.** Evaluation of oregano essential oil (*Origanum heracleoticum* L.) on growth, antioxidant effect and resistance against *Aeromonas hydrophila* in channel catfish (*Ictalurus punctatus*). *Aquaculture*, 292(3-4), pp.214-218.
- Zhou, L., Zheng, H., Tang, Y., Yu, W. and Gong, Q., 2013.** Eugenol inhibits quorum sensing at sub-inhibitory concentrations. *Biotechnology Letters*, 35(4), pp.631-637.
- Zhu, X., Davis, D.A., Roy, L.A., Samocha, T.M. and Lazo, J.P., 2013a.** Response of Pacific white shrimp, *Litopenaeus vannamei*, to three sources of solvent extracted soybean meal. *Journal of the World Aquaculture Society*, 44(3), pp.396-404.
- Zhou, E., Fu, Y., Wei, Z., Yu, Y., Zhang, X. and Yang, Z., 2014.** Thymol attenuates allergic airway inflammation in ovalbumin (OVA)-induced mouse asthma. *Fitoterapia*, 96, pp.131-137.
- Zhou, S., Li, W.X., Wang, Y.Q., Zou, H., Wu, S.G. and Wang, G.T., 2017.** Anthelmintic efficacies of three common disinfectants and extracts of four traditional Chinese medicinal plants against *Gyrodactylus kobayashii* (Monogenea) in goldfish (*Carassius auratus*). *Aquaculture*, 466, pp.72-77.
- Zilberg, D., Tal, A., Froyman, N., Abutbul, S., Dudai, N. and Golan-Goldhirsh, A., 2010.** Dried leaves of *Rosmarinus officinalis* as a treatment for streptococcosis in tilapia. *Journal of Fish Diseases*, 33(4), pp.361-369.